



# Sickle cell disorder in adulthood

## Patient information factsheet

### Introduction

- This leaflet explains more about sickle cell disorder (SCD) in adults and provides an overview of the services offered by Whittington Health.
- If you would like further information, please speak to a member of our team.

### What is SCD?

- SCD is a disorder of the body's red blood cells that is genetically inherited and lifelong.
- It is caused by an abnormality in haemoglobin (the part of red blood cells that carries oxygen round the body), which makes the red cells become inflexible and sticky. This in turn makes other blood cells, as well as the vessels they travel in, sticky.
- Blood flow becomes interrupted, and an inflammatory cascade occurs – a process where one inflammatory reaction causes another.
- This process may cause pain and problems in the organs where the sickling is happening.
- It also affects the immune system.
- In SCD, there are problems that happen suddenly and are often painful – these are known as sickle cell crises.
- The long-term effects of crises and background sickling (that may have no symptoms) are known as chronic complications.
- The approach we take is to limit and prevent both of these from happening as much as possible.

### What kind of problems can happen in SCD?

- Infection
- Gallstones
- Joint damage/destruction
- Loss of spleen function (making patients prone to infections)
- Damage to kidneys
- Leg ulcers
- Visual impairment
- Neurological damage which may lead to stroke
- Erectile dysfunction (priapism).



## The signs and symptoms of crisis may include:

- Fatigue (feeling tired or weak)
- Pain
- Jaundice (yellowing of the white of the eyes)
- Paleness of skin, or inside the mouth or eyes
- Shortness of breath
- Dizziness
- Headaches
- Symptoms of infection, such as fever
- **If you develop any of the following symptoms: high fever, chest pain or pain that does not feel like the usual sickle pain, shortness of breath, weakness (especially one sided), change in vision or extreme tiredness, please seek medical attention straight away.**

## Treatment

- You may be able to manage a simple painful sickle cell crisis at home by taking painkillers, resting, drinking fluids and keeping warm.
- However, if the pain becomes worse you may need stronger painkillers in hospital.
- We can help to control the pain in hospital as well as treat the underlying cause (trigger) and any complications relating to your condition.
- Sometimes the crisis may not be particularly painful or more importantly, you may feel unwell from the start. **If you are unwell, even if you have no pain, you should always contact us as you may need to come to hospital.**
- **We may also suggest other ways of managing your condition with preventative treatments.** We usually recommend this if the frequency of sickle cell crises increases or if an attack was particularly severe. **Treatment options include starting hydroxyurea or a long-term blood transfusion programme.**
- **Some people have had a bone marrow transplant that has cured their sickle cell disease.**

## What to do in an emergency

- If you have a non-urgent medical problem or a medical problem unrelated to your sickle cell disease, for example a rash or a twisted ankle, you can contact your GP or go to your local Emergency Department (A&E). If necessary, they can discuss your problem with us.
- If you have a non-urgent problem related to sickle cell disease, you can ask for your outpatient appointment to be brought forward.
- **If you are more unwell beyond your usual sickle cell crisis, you should attend the Whittington Hospital Emergency Department (A&E).**



- **If you attend an Emergency Department (A&E) in another hospital, please inform us so we can liaise with the doctors looking after you in that hospital.**

## What is an emergency?

The following symptoms require **immediate medical attention**:

- Fever: 38°C or above
- Feeling very unwell
- Pain that cannot be controlled with painkillers at home
- New weakness, particularly if felt more on one side than the other
- If you think you are much more anaemic than usual
- Shortness of breath with or without cough
- Tummy problems – severe pain, diarrhoea, vomiting
- Priapism: a prolonged and painful erection that happens without sexual arousal or stimulation, and can last for more than four hours (see Priapism information leaflet)

## What can trigger a sickle cell crisis?

- Common triggers of a sickle cell crisis include infection, stress, dehydration, cold/hot weather or sudden changes in temperature, or nothing at all.
- Sometimes crises happen out of the blue.

## How to prevent SCD

- As SCD is an inherited condition, each carrier (trait) parent may pass on an affected gene to their child.
- If both parents are carriers, the chance that their child will have SCD is one in four.
- If one person has the disease and the other is a carrier, the risk is greater at one in two.
- Therefore, it is important that your partner is tested for SCD if you are planning a pregnancy.
- **If you would like to discuss this further, please speak to us or go to your local sickle cell counselling service.**
- If you or your partner becomes pregnant, it is extremely important that testing is done as early as possible. Please contact us immediately if this happens and we will organise further counselling and assessment.



## What can I do to stay well?

- Maintain a good fluid intake
- Eat healthily
- Exercise
- Make sure infections are treated quickly
- Maintain a good balance of nutrition and activity
- Ensure your vaccinations are up to date
- Avoid smoking
- Take penicillin twice a day for life
- Ensure that you attend your clinic appointments so that we can review your health and monitor you for any complications.
- **Attending your clinic appointments is a recommendation stipulated in the National Standards produced by the Sickle Cell Society together with the Department of Health. These appointments are every six to 12 months in healthy people but may be more frequent if you have health problems.**
- **If you do not have an appointment, please ask your GP to refer you.**
- If your appointment is not convenient, please change it. We understand that you have a busy life with many commitments, and we will do our best to schedule an appointment that is convenient for you.
- For further information on how to stay well, please see our leaflet: Medications, vaccinations and travel in sickle cell disease.

## Contact details

- Your consultant team is:

Dr Emma Drasar  
 Dr Ryan Mullally  
 Dr Ana Ortuno-Cabrero  
 Dr Ali Rismeni  
 Dr Zara Sayar  
 Dr Annabel McMillan  
 Dr Prabal Mittal  
 Dr Janine Younis



- Adult departmental contacts:  
[whh-tr.haematologyteam@nhs.net](mailto:whh-tr.haematologyteam@nhs.net) Haematology Administration Team  
[whh-tr.adultsicklecons@nhs.net](mailto:whh-tr.adultsicklecons@nhs.net) Adult Sickle Cell Clinical Nurse Specialist (CNS) team  
[ncl.redcellteam@nhs.net](mailto:ncl.redcellteam@nhs.net) Community link (Sickle cell, Thalassaemia and Rare Anaemia)  
 0203 316 8853 Community phone number  
[emma.prescott@nhs.net](mailto:emma.prescott@nhs.net) Whittington Thalassaemia CNS  
[niamh.malone-cooke@nhs.net](mailto:niamh.malone-cooke@nhs.net) Whittington Thalassaemia CNS
- Network website:  
<https://www.uclh.nhs.uk/theredcellnetwork/subscribe>

If you or your family have any other questions, please do not hesitate to contact any of the above medical team at Whittington Health.

## Where can I get more information?

### The Sickle Cell Society

54 Station Rd, London NW10 4UA

Tel: 020 8861 7795

Website: <http://www.sicklecellsociety.org>

### Red Cells R Us

Sickle Cell & Thalassaemia Centre

17a Hornsey Street, London N7 8GG

Tel: 020 3316 8853/8854

Twitter: @Red Cells R Us

Facebook: Red Cells R Us

Google+: Red Cells R Us (in community section)

### NHS 111

Tel: 111

[www.nhs.uk/NHSEngland/AboutNHSservices/Emergencyandurgentcareservices/Pages/NHS-111.aspx](http://www.nhs.uk/NHSEngland/AboutNHSservices/Emergencyandurgentcareservices/Pages/NHS-111.aspx)



Patient advice and liaison service (PALS)

If you have a compliment, complaint or concern please contact our PALS team on 020 7288 5551 or [whh-tr.PALS@nhs.net](mailto:whh-tr.PALS@nhs.net)

If you need a large print, audio or translated copy of this leaflet please email [whh-tr.patient-information@nhs.net](mailto:whh-tr.patient-information@nhs.net). We will try our best to meet your needs.

Twitter.com/WhitHealth  
Facebook.com/WhittingtonHealth

Whittington Health NHS Trust  
Magdala Avenue  
London  
N19 5NF  
Phone: 020 7272 3070  
[www.whittington.nhs.uk](http://www.whittington.nhs.uk)

Date published: 25/09/2024  
Review date: 25/09/2026  
Ref: EIM/Haem/SCDAAd/02

© Whittington Health  
Please recycle

